



34123

VCN Supplement

Part D

Specifications

Appearance (Colour)	White, fine powder
Solubility	Soluble in distilled water
Cultural Response	Inoculate and incubate at $35 \pm 2^\circ\text{C}$ for 18-48 hours (Aseptically add the contents of one vial of VCN Supplement (34123) as per direction)
Organism	Neisseria meningitidis ATCC 13090
Inoculum (cfu) 10-100	Growth : Good Colony colour : Medium to large, blue-gray, mucoid
Organism	Escherichia coli ATCC 25922
Inoculum	Growth : Inhibited
Organism	Proteus mirabilis ATCC 25933
Inoculum	Growth : Inhibited

Other Information

Applications

Used for selective isolation of Neisseria species.

Composition

Vancomycin	1.50 mg/vial
Colistin methane sulphonate	3.75 mg/vial
Nystatin	6250 units/vial

Directions

1. Aseptically add 2 ml of sterile distilled water to 1 vial.
2. Mix well and aseptically add to sterile cooled media at $45-50^\circ\text{C}$ and dispense as desired.
3. Add 2 ml (1 vial) of VCN supplement to 440 ml of Thayer-Martin Medium, Base (44521), Transgrow Medium.
4. Add 2 ml (1 vial) of VCN supplement to 250 ml of GC Agar Base(72464).

General Information

Storage	2 to 8°C (Refrigerate)
Shelf Life	24 Months
IMDG Identification	Not Regulated for Transport (Non-Haz)
HSN Code	
5 vl	38229090 (GST 12%)
5 x5 vl	38229090 (GST 12%)

Available Packages

5 vI

5 x5 vI

Disclaimer

The information represented here may/may not represent the entire product specification, application or protocol recommended by Sisco Research Laboratories Pvt. Ltd. (SRL). This information is for the user scientists or trading community as a guide in their applications. The company claims no liability for misuse resulting due to wrong usage of the information above. For actual batch related documents, mail us.

608-B, Satellite Gazebo, Andheri Ghatkopar Link Road, Chakala, Andheri (E), Mumbai - 400 099, Maharashtra, India. Telephone: +91-22-4268 5800, Email: info@srchem.com, website www.srchem.com
