



47082

## Enterococcus Selective Supplement

Part D

### Specifications

Appearance (Colour)	White, fine powder
Solubility	Soluble in 0.2 N Sodium hydroxide
Cultural Response	Inoculate and incubate at $44 \pm 2^\circ\text{C}$ for 36-72 hours. (Aseptically add the rehydrated contents of 1 vial as per direction) Growth observed on MUD SF Broth.
Organism	Enterococcus faecalis ATCC 29212
Inoculum (cfu) 10-100	Growth : Good Flourescence at 366 nm : Positive (Blue)

### Other Information

#### Applications

Used for the selective isolation and cultivation of Enterococci.

#### Composition

Nalidixic Acid	25.00 mg/vial
Thallium acetate	200.00 mg/vial

#### Directions

1. Aseptically add 5 ml of sterile 0.2N sodium hydroxide to 1 vial.
2. Mix well and aseptially add sterile cooled media at  $40-45^\circ\text{C}$  and pour in petriplates.
3. Add 5ml (1 vial) of Entreococcus selective supplement to 100 ml of MUD SF Broth alongwith sterile 1% TTC solution.

### General Information

Storage	2 to $8^\circ\text{C}$ (Refrigerate)
Shelf Life	24 Months
IMDG Identification	Not Regulated for Transport (Non-Haz)
HSN Code	
5 vl	38229090 (GST 12%)
5 x5 vl	38229090 (GST 12%)

### Available Packages

5 vl

5 x5 vl

#### Disclaimer

The information represented here may/may not represent the entire product specification, application or protocol recommended by Sisco Research Laboratories Pvt. Ltd. (SRL). This information is for the user scientists or trading community as a guide in their applications. The company claims no liability for misuse resulting due to wrong usage of the information above. For actual batch related documents, mail us.

608-B, Satellite Gazebo, Andheri Ghatkopar Link Road, Chakala, Andheri (E), Mumbai - 400 099, Maharashtra, India. Telephone: +91-22-4268 5800, Email: info@srchem.com, website www.srchem.com