

91338

Potato Dextrose Agar (PDA) BioVeg

Part D

Specifications	
Appearance (Colour)	Light cream
Appearance (Form)	Free flowing, h
Solubility	39.00 gm/liter
Solubility before autoclaving (Clarity)	Very slightly op
Gel strength (1.5% gel)	Firm, comparab
pH (25°C)	5.6 ± 0.2
Prepared Medium Appearance after autoclaving (Clarity)	Very slightly opa
Prepared Medium Appearance after autoclaving (Colour)	Light amber
Cultural Response	Inoculate and in 18-48 hours or u
Organism	Candida albican
Inoculum (cfu) 10-100	Growth: Good
Organism	Saccharomyces
Inoculum (cfu) 10-100	Growth: Good
Organism	Aspergillus niger
Inoculum	Growth : Good

Other Information

Applications

For the isolation, cultivation and enumeration of yeasts and molds from dairy and food products. Used to induce sporulation in many fungi.

Composition	
Ingredients	gm/lt.
Potatoes infusion from (Veg)	200.00
Dextrose	20.00
Agar	15.00

Directions

- 1. Add 39.00 gm powder to distilled/purified water.
- 2. Bring volume to 1.0 litre and mix thoroughly.
- 3. Gently heat and bring to boiling.
- 4. Autoclave at 15 psi pressure at 121°C for 15 minutes.
- 5. Dispense into sterile petri plates.

General Information

Storage	8 to 25°C (Cool & Dry Area)
Shelf Life	36 Months
IMDG Identification	Not Regulated for Transport (Non-Haz)

HSN Code	
100 Gms	38210000 (GST 18%)
500 Gms	38210000 (GST 18%)

Available Packages

100 Gms

500 Gms

Disclaimer